

A REVISED CENSUS OF THE TERRESTRIAL
MOLLUSCA OF TASMANIA.

By W. F. PETTERD (Launceston) and C. HEDLEY, Conchologist.

(Plates lxxxii.-lxxxvii, and fig. 16.)

This paper is confined to the rehabilitation of species. We have endeavoured to supply illustrations of all Tasmanian land shells hitherto unfigured, to simplify the tangled synonymy and to arrange the species in genera and higher groups in harmony with modern classification.

Our work has been facilitated by the loan of the collection of the Tasmanian Museum, for which we are indebted to the kindness of the late Mr. Alexander Morton, the then Curator. Examples of most of the species have been added to the Australian Museum Collection.

Group O R T H U R E T H R A.

Family PUPIDÆ.

Genus V E R T I G O, Müller, 1774.

V E R T I G O L I N C O L N E N S I S, Cox.

Pupa lincolnensis, Cox, Proc. Zool. Soc., 1867, p. 39; *Id.*, Mon. Austr. Land Shells, 1868, p. 80, pl. xiv., f. 16. *Id.*, Pilsbry, Proc. Acad. Nat. Sci. Philad., 1900, p. 428. *Id.*, (*P. tasmanica*) Johnston, Proc. Roy. Soc. Tas., 1882 (1883) p. 144, pl. i.

Hab., Swansea. Obtained plentifully on the sand-dunes; extends to South Australia.

Group H E T E R U R E T H R A.

Family SUCCINEIDÆ.

Genus S U C C I N E A, Draparnaud, 1801.

S U C C I N E A A U S T R A L I S, Ferrussac.

Succinea australis, Ferrussac, Tabl. Syst., ii., 1821, p. 27. *Id.*, Gray, Ann. Phil., 1825, p. 415, pl. ix. *Id.*, Ferrussac and Deshayes, Hist. Nat. Moll. Terr. et Fluv., ii., pt. 2, n.d., p. 137, pl. xi., fig. 11. *Id.*, Quoy et Gaim. Voy. Astrolabe, Zool., ii., 1832, p. 150, pl. xiii., f. 19-23.

S. legrandi, Cox in Legrand, Coll. Mon., 1871, sp. 2.